

Lemma 1.11 For any ordering of the grid points, the matrix A of the system (1.6) is symmetric and negative definite.

Proof. Equation (1.6) implies that if $a_{ij} \neq 0$ for $i \neq j$, then the i -th and j -th points of the grid (ph, qh) , are nearest neighbours. Hence $a_{ij} \neq 0$ implies $a_{ji} = a_{ji} = 1$, which proves the symmetry of A . Therefore A has real eigenvalues and eigenvectors.

It remains to prove that all the eigenvalues are negative. The arguments are parallel to the proof of Gershgorin theorem. Let $Ax = \lambda x$, and let i be an integer such that $|x_i| = \max |x_j|$. With such an i we address the following identity (which is a reordering of the equation $(Ax)_i = \lambda x_i$):

$$\underbrace{|(\lambda - a_{ii})x_i|}_{|\lambda+4||x_i|} = \underbrace{|\sum_{j \neq i}^n a_{ij}x_j|}_{\leq 4|x_i|} \quad (1.7)$$

Here $a_{ii} = -4$ and $a_{ij} \in \{0, 1\}$ for $j \neq i$, with at most four nonzero elements on the right-hand side. It is seen that the case $\lambda > 0$ is impossible. Assuming $\lambda = 0$, we obtain $|x_j| = |x_i|$ whenever $a_{ij} = 1$, so we can alter the value of i in (1.7) to any of such j and repeat the same arguments. Thus, the modulus of every component of x would be $|x_i|$, but then the equations (1.7) that occur at the boundary of the grid and have fewer than four off-diagonal terms (see (1.6)) could not be true. Hence, $\lambda = 0$ is impossible too, hence $\lambda < 0$ which proves that A is negative definite. \square

Proposition 1.12 The eigenvalues of the matrix A are

$$\lambda_{k,\ell} = -4 \left(\sin^2 \frac{k\pi h}{2} + \sin^2 \frac{\ell\pi h}{2} \right), \quad h = \frac{1}{m+1}, \quad k, \ell = 1 \dots m,$$

Proof. Let us show that, for every pair (k, ℓ) , the vectors

$$v = (v_{i,j}), \quad v_{i,j} = \sin ix \sin jy, \quad \text{where } x = k\pi h, \quad y = \ell\pi h,$$

are the eigenvectors of A . Indeed, for $i, j = 1 \dots m$, we have

$$\begin{aligned} (Av)_{i,j} &= \sin(jy) [\sin(ix - x) - 2\sin(ix) + \sin(ix + x)] \\ &\quad + \sin(ix) [\sin(jy - y) - 2\sin(jy) + \sin(jy + y)] \\ &= \sin(jy) \sin(ix) [2\cos x - 2] + \sin(ix) \sin(jy) [2\cos y - 2] = \lambda v_{i,j}. \end{aligned}$$

Note that the terms $u_{i \pm 1, j}$, $u_{i, j \pm 1}$ do not appear in (1.6) for $i, j = 1$ or $i, j = m$, respectively, therefore (for such i, j) we should have dropped the corresponding components from above equation, but they are equal to zero because $\sin(i-1)x = 0$ for $i = 1$, while $\sin(i+1)x = 0$ for $i = m$, since $x = \frac{k\pi}{m+1}$. Thus, the eigenvalues are

$$\lambda_{k,\ell} = [2\cos x - 2] + [2\cos y - 2] = -4 \left(\sin^2 \frac{x}{2} + \sin^2 \frac{y}{2} \right) = -4 \left(\sin^2 \frac{k\pi h}{2} + \sin^2 \frac{\ell\pi h}{2} \right). \quad \square$$

Remark 1.13 As a matter of independent mathematical interest, note that for $1 \leq k, \ell \ll m$ we have $\sin x \approx x$, hence the eigenvalues for the discretized Laplacian ∇_h^2 are

$$\frac{\lambda_{k,\ell}}{h^2} \approx -\frac{4}{h^2} \left[\frac{k^2\pi^2 h^2}{4} + \frac{\ell^2\pi^2 h^2}{4} \right] = -(k^2 + \ell^2)\pi^2.$$

Now, recall (e.g. from the solution of the Poisson equation in a square by separation of variables in Maths Methods) that the *exact* eigenvalues of ∇^2 (in the unit square) are $-(k^2 + \ell^2)\pi^2$, $k, \ell \in \mathbb{N}$, with the corresponding eigenfunctions $V_{k,\ell}(x, y) = \sin k\pi x \sin \ell\pi y$. So, the eigenvectors of the discretized ∇_h^2 are the values of $V_{k,\ell}(x, y)$ on the grid-points, and the eigenvalues of ∇_h^2 approximate those for continuous case.