

Mathematical Tripos Part II: Michaelmas Term 2014

Numerical Analysis – Lecture 9

Problem 2.29 (The diffusion equation in two space dimensions) We are solving

$$\frac{\partial u}{\partial t} = \nabla^2 u, \quad 0 \leq x, y \leq 1, \quad t \geq 0, \quad (2.14)$$

where $u = u(x, y, t)$, together with initial conditions at $t = 0$ and Dirichlet boundary conditions at $\partial\Omega$, where $\Omega = [0, 1]^2 \times [0, \infty)$. It is straightforward to generalize our derivation of numerical algorithms, e.g. by the method of lines. Thus, let $u_{\ell, m}(t) \approx u(\ell h, m h, t)$, where $h = \Delta x = \Delta y$, and let $u_{\ell, m}^n \approx u_{\ell, m}(n h)$ where $k = \Delta t$. The five-point formula results in

$$u'_{\ell, m} = \frac{1}{h^2} (u_{\ell-1, m} + u_{\ell+1, m} + u_{\ell, m-1} + u_{\ell, m+1} - 4u_{\ell, m}),$$

or in the matrix form

$$\mathbf{u}' = \frac{1}{h^2} A_* \mathbf{u}, \quad \mathbf{u} = (u_{\ell, m}) \in \mathbb{R}^N, \quad (2.15)$$

where A_* is the block TST matrix of the five-point scheme:

$$A_* = \begin{bmatrix} H & I & & & \\ & I & \ddots & & \\ & & \ddots & \ddots & \\ & & & I & H \end{bmatrix}, \quad H = \begin{bmatrix} -4 & 1 & & & \\ & 1 & \ddots & & \\ & & \ddots & \ddots & \\ & & & 1 & \\ & & & & 1 & -4 \end{bmatrix}.$$

Thus, the Euler method yields

$$u_{\ell, m}^{n+1} = u_{\ell, m}^n + \mu (u_{\ell-1, m}^n + u_{\ell+1, m}^n + u_{\ell, m-1}^n + u_{\ell, m+1}^n - 4u_{\ell, m}^n), \quad (2.16)$$

or in the matrix form

$$\mathbf{u}^{n+1} = A \mathbf{u}^n, \quad A = I + \mu A_*$$

where, as before, $\mu = \frac{k}{h^2} = \frac{\Delta t}{(\Delta x)^2}$. The local error is $\eta = \mathcal{O}(k^2 + k h^2) = \mathcal{O}(h^4)$. To analyse stability, we notice that A is symmetric, hence normal, and its eigenvalues are related to those of A_* by the rule

$$\lambda_{k, \ell}(A) = 1 + \mu \lambda_{k, \ell}(A_*) \quad \underbrace{=} \quad 1 - 4\mu \left(\sin^2 \frac{\pi k h}{2} + \sin^2 \frac{\pi \ell h}{2} \right).$$

Proposition 1.12

Consequently,

$$\sup_{h>0} \rho(A) = \max\{1, |1 - 8\mu|\}, \quad \text{hence} \quad \mu \leq \frac{1}{4} \Leftrightarrow \text{stability}.$$

Method 2.30 (Fourier analysis) Fourier analysis generalizes to two dimensions: of course, we now need to extend the range of (x, y) in (2.14) from $0 \leq x, y \leq 1$ to $x, y \in \mathbb{R}$. A 2D Fourier transform reads

$$\widehat{u}(\theta, \psi) = \sum_{\ell, m \in \mathbb{Z}} u_{\ell, m} e^{-i(\ell\theta + m\psi)}$$

and all our results readily generalize. In particular, the Fourier transform is an isometry from $\ell_2[\mathbb{Z}^2]$ to $L_2([-\pi, \pi]^2)$, i.e.

$$\left(\sum_{\ell, m \in \mathbb{Z}} |u_{\ell, m}|^2 \right)^{1/2} =: \|\mathbf{u}\| = \|\widehat{u}\|_* := \left(\frac{1}{4\pi^2} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} |\widehat{u}(\theta, \psi)|^2 d\theta d\psi \right)^{1/2},$$

and the method is stable iff $|H(\theta, \psi)| \leq 1$ for all $\theta, \psi \in [-\pi, \pi]$. The proofs are an easy elaboration on the one-dimensional theory. Insofar as the Euler method (2.16) is concerned,

$$H(\theta, \psi) = 1 + \mu (e^{-i\theta} + e^{i\theta} + e^{-i\psi} + e^{i\psi} - 4) = 1 - 4\mu \left(\sin^2 \frac{\theta}{2} + \sin^2 \frac{\psi}{2} \right),$$

and we again deduce stability if and only if $\mu \leq \frac{1}{4}$.

Method 2.31 (Crank-Nicolson for 2D) Applying the trapezoidal rule to our semi-dcretization (2.15) we obtain the two-dimensional Crank-Nicolson method:

$$(I - \frac{1}{2}\mu A_*) \mathbf{u}^{n+1} = (I + \frac{1}{2}\mu A_*) \mathbf{u}^n, \quad (2.17)$$

in which we move from the n -th to the $(n+1)$ -st level by solving the system of linear equations $B\mathbf{u}^{n+1} = C\mathbf{u}^n$, or $\mathbf{u}^{n+1} = B^{-1}C\mathbf{u}^n$. For stability, similarly to the one-dimensional case, the eigenvalue analysis implies that $A = B^{-1}C$ is normal and shares the same eigenvectors with B and C , hence

$$\lambda(A) = \frac{\lambda(C)}{\lambda(B)} = \frac{1 + \frac{1}{2}\mu\lambda(A_*)}{1 - \frac{1}{2}\mu\lambda(A_*)} \Rightarrow |\lambda(A)| < 1 \text{ as } \lambda(A_*) < 0$$

and the method is stable for all μ . The same result can be obtained through the Fourier analysis.

Matlab demo: Download the Matlab GUI for *Solving the Wave and Diffusion Equations in 2D* from http://www.maths.cam.ac.uk/undergrad/course/na/ii/pdes_2d/pdes_2d.php and solve the diffusion equation (2.14) for different initial conditions. For the numerical solution of the equation you can choose from the Euler method and the Crank-Nicolson scheme. The GUI allows you to solve the wave equation as well. Compare the behaviour of solutions!

Technique 2.32 (Splitting) We would like to find a fast solver to the system (2.17). The matrix $B = I - \frac{1}{2}\mu A_*$ has a structure similar to that of A_* , so we may apply the Hockney method. However, since the method (2.17) has a local truncation error $\mathcal{O}(k^3 + kh^2)$, we don't need an exact solution of the system: it would be enough to have one within the error.

Let us employ the notation

$$\Delta_x^2 u_{\ell,m} = u_{\ell-1,m} - 2u_{\ell,m} + u_{\ell+1,m}, \quad \Delta_y^2 u_{\ell,m} = u_{\ell,m-1} - 2u_{\ell,m} + u_{\ell,m+1}.$$

Then the Crank-Nicolson method calculates \mathbf{u}^{n+1} by solving the system

$$[I - \frac{1}{2}\mu(\Delta_x^2 + \Delta_y^2)] u_{\ell,m}^{n+1} = [I + \frac{1}{2}\mu(\Delta_x^2 + \Delta_y^2)] u_{\ell,m}^n, \quad \ell, m = 1 \dots M. \quad (2.18)$$

The local error is however preserved if we replace this formula by the difference equation

$$[I - \frac{1}{2}\mu\Delta_x^2][I - \frac{1}{2}\mu\Delta_y^2] u_{\ell,m}^{n+1} = [I + \frac{1}{2}\mu\Delta_x^2][I + \frac{1}{2}\mu\Delta_y^2] u_{\ell,m}^n, \quad (2.19)$$

which is called the split version of Crank-Nicolson. Indeed, the difference between two schemes is equal to

$$\begin{aligned} \frac{1}{4}\mu^2\Delta_x^2\Delta_y^2(u_{\ell,m}^{n+1} - u_{\ell,m}^n) &= \frac{k^2}{4}\frac{1}{h^2}\Delta_x^2\frac{1}{h^2}\Delta_y^2\left(k\frac{\partial}{\partial t}u_{\ell,m}^n + \mathcal{O}(k^2)\right) \\ &= \frac{k^3}{4}\left(\frac{\partial^2}{\partial x^2}\frac{\partial^2}{\partial y^2}\frac{\partial}{\partial t}u_{\ell,m}^n + \mathcal{O}(k+h^2)\right) = \mathcal{O}(k^3 + kh^2), \end{aligned}$$

the same magnitude as of the local error. In the matrix form, (2.19) is equivalent to splitting the matrix A_* into the sum of two matrices A_x and A_y as

$$A_* = A_x + A_y, \quad A_x = \begin{bmatrix} -2I & I & & & \\ & I & \ddots & & \\ & & \ddots & \ddots & \\ & & & \ddots & I \\ & & & & I & -2I \end{bmatrix}, \quad A_y = \begin{bmatrix} H & & & & \\ & H & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & H \end{bmatrix}, \quad H = \begin{bmatrix} -2 & 1 & & & \\ & 1 & \ddots & & \\ & & \ddots & \ddots & \\ & & & \ddots & 1 \\ & & & & 1 & -2 \end{bmatrix}$$

and solving the uncoupled system

$$[I - \frac{1}{2}\mu A_x][I - \frac{1}{2}\mu A_y] \mathbf{u}^{n+1} = [I + \frac{1}{2}\mu A_x][I + \frac{1}{2}\mu A_y] \mathbf{u}^n.$$

as

$$B_x \mathbf{u}^{n+1/2} = C_x C_y \mathbf{u}^n, \quad B_y \mathbf{u}^{n+1} = \mathbf{u}^{n+1/2}.$$

Matrix $B_y = I - \frac{1}{2}\mu A_y$ is block diagonal, and solving $B_y \mathbf{u} = \mathbf{v}$ is just solving one and the same tridiagonal system $B\mathbf{u}_i = \mathbf{v}_i$ with different right-hand sides. Matrix $B_x = I - \frac{1}{2}\mu A_x$ is of the same form up to a permutation (reordering of the grid), so solving $B_x \mathbf{v} = \mathbf{b}$ is again a fast procedure.